THE JOURNAL OF CHEMICAL THERMODYNAMICS

Volume 17, 1985

EDITORS

M. L. McGlashan
Department of Chemistry
University College London
20 Gordon Street
London WC1H 0AJ
U.K.

P. A. G. O'Hare
Argonne National Laboratory
Chemical Technology Division
Argonne
Illinois 60439
U.S.A.

Assistant Editor: **M. B. Ewing** (University College London, U.K.)

ADVISORY BOARD

Chairman: G. M. Schneider (F.R.G.)

S. Angus (U.K.)
G. C. Benson (Canada)
J. C. G. Calado (Portugal)
M. Colomina (Spain)
J. D. Cox (U.K.)
M. Gaune-Escard (France)
C. Jolicoeur (Canada)
O. J. Kleppa (U.S.A.)
M. Månsson (Sweden)
V. A. Medvedev (U.S.S.R.)
F. L. Oetting (U.S.A.)
J. B. Ott (U.S.A.)
F. Pavese (Italy)
G. Somsen (The Netherlands)
J. Stecki (Poland)
H. Suga (Japan)
F. L. Swinton (U.K.)
A. G. Williamson (N.Z.)

ACADEMIC PRESS
London Orlando San Diego
San Francisco New York Toronto
Montreal Sydney Tokyo São Paulo
A Subsidiary of Harcourt Brace Jovanovich, Publishers

Copyright © 1985 Academic Press Inc. (London) Limited

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 17

Number 1, January 1985

CHRISTENSEN, J. J.; ZEBOLSKY, D. M.; IZATT, R. M. The excess enthalpies of (carbon dioxide + toluene) at 470.15 and 573.15 K from 7.60 to 12.67 MPa 1

BEYER, R. P.; BENNINGTON, K. O.; BROWN, R. R. Sodium zirconium oxide: molar enthalpy of formation at 298.15 K and molar heat capacity from 5 to 1168 K 11

CORDFUNKE, E. H. P.; OUWELTJES, W.; PRINS, G. Standard enthalpies of formation of uranium compounds. XI. Lithium uranates(VI) . . . 19

RODRIGUEZ-NUÑEZ, E.; PAZ-ANDRADE, M. I.; JIMENEZ, E.; BRAVO, R. Thermodynamic properties of (a xylene + an *n*-alkanol). II. Excess molar enthalpies at 298.15 K for 1-propanol, 1-butanol, and 1-pentanol . . 23

MARSH, K. N. Excess enthalpies and excess volumes of nitromethane +, and nitroethane + each of several non-polar liquids 29

PASHINKIN, A. S.; RABINOVICH, I. B.; SHEIMAN, M. S.; NISTRATOV, V. P.; VOROBJOVA, O. I. Heat capacity and thermodynamic functions of tellurium dioxide in the range 5 to 300 K 43

EASTEAL, A. J.; WOOLF, L. A. (p, V_m, T, x) measurements for $\{(1-x)H_2O + xCH_3OH\}$ in the range 278 to 323 K and 0.1 to 280 MPa. I. Experimental results, isothermal compressibilities, thermal expansivities, and partial molar volumes 49

GUARIDO, C. G.; SUAREZ, M.; GARCIA, J. R.; LLAVONA, R.; RODRIGUEZ, J. Thermodynamic treatment and calorimetric study of H^+/Li^+ ion exchange on α-titanium phosphate 63

EASTEAL, A. J.; WOOLF, L. A. (p, V_m, T, x) measurements for $\{(1-x)H_2O + xCH_3OH\}$ in the range 278 to 323 K and 0.1 to 280 MPa. II. Thermodynamic excess properties 69

BOTTINI, S. B.; SAVILLE, G. Excess enthalpies for (water + nitrogen)(g) and (water + carbon dioxide)(g) at 520 to 620 K and up to 4.5 MPa . . 83

NOTE

ARNOLD, P.; LILLEY, T. H. Aqueous solutions containing amino acids and peptides. 14. The enthalpy of interaction of β-alanine and urea . . 99

Number 2, February 1985

CHAUDHARI, S. K.; KATTI, S. S. Excess volumes of (an isomer of butanol + *n*-octane) at 298.15 K, measured with a continuous-dilution dilatometer 101

AWWAD, A. M.; AL-MADFAI, S. F.; JBARA, K. A.; KANBOUR, F. I. Excess volumes of (an aromatic hydrocarbon + *N*-formylmorpholine) . . 105

TOPOR, L.; KLEPPA, O. J. Molar enthalpy of formation of CrB_2 by high-temperature calorimetry 109

RAVIPRASAD, A.; VENKATESWARA RAO, K. Vapour pressure of 2,4-pentanedione 117

TRESZCZANOWICZ, A. J.; BENSON, G. C. Excess volumes for (2-methylbutan-2-ol + *n*-heptane) and for (cyclopentanol + *n*-heptane) . . 123

BAIDAKOV, V. G.; GURINA, T. A. (p, ρ, T) of superheated liquid oxygen . 131

SHEALY, G. S.; SANDLER, S. I. Vapor–liquid equilibria of $\{xCH_3CH(CH_3)CHO + (1-x)C_7H_{16}\}$(l) and $\{xCH_3CO_2C_2H_5 + (1-x)C_7H_{16}\}$(l) 143

KRATZKE, H.; MÜLLER, S. Thermodynamic properties of acetonitrile. 2. (p, ρ, T) of saturated and compressed liquid acetonitrile 151

MASUDA, Y.; MIYAMOTO, H.; KANEKO, Y.; HIROSAWA, K. The standard molar enthalpies of formation of crystalline rubidium and cesium oxalates . 159

ROBIE, R. A.; HEMINGWAY, B. S. Low-temperature molar heat capacities and entropies of MnO_2 (pyrolusite), Mn_3O_4 (hausmanite), and Mn_2O_3 (bixbyite) 165

CHRISTENSEN, J. J.; ZEBOLSKY, D. M.; IZATT, R. M. The excess enthalpies of (carbon dioxide + hexane) at 470.15, 510.15, and 573.15 K from 7.50 to 12.50 MPa 183

DETHLEFSEN, C.; HVIDT, AA. Densities and derived volume functions of binary mixtures: (an ethylene glycol derivative + water) at 298.15 K . . 193

Number 3, March 1985

HANDA, Y. P. Heat capacities in the range 95 to 260 K and enthalpies of fusion for structure-II clathrate hydrates of some cyclic ethers . . 201

HAYNES, W. M.; MCCARTY, R. D.; EATON, B. E.; HOLSTE, J. C. Isochoric (p, V_m, x, T) measurements on (methane + ethane) from 100 to 320 K at pressures to 35 MPa 209

KANA'AN, A. S. Thermodynamics of the sublimation of thallium(I) fluoride by simultaneous Knudsen-effusion and torsion-effusion 233

MARUSHKIN, K. N.; ALIKHANYAN, A. S.; GREENBERG, J. H.; LAZAREV, V. B.; MALYUSOV, V. A.; ROZANOVA, O. N.; MELEKH, B. T.; GORGORAKI, V. I. Sublimation thermodynamics of tungsten trioxide 245

WIECZOREK, S. A.; SIPOWSKA, J. T. Vapour pressures and excess Gibbs free energies of (2,3,3-trimethylbutan-2-ol + *n*-heptane) between 298.150 and 363.279 K 255

DRAEGER, J. A. The methylbenzenes. II. Fundamental vibrational shifts, statistical thermodynamic functions, and properties of formation . . 263

CARSON, A. S.; LAYE, P. G.; SPENCER, J. A. The enthalpies of formation of triphenyl vinyl tin and triphenyl phenylethynyl tin and some associated tin–carbon bond enthalpies 277

KRATZKE, H.; MÜLLER, S.; BOHN, M.; KOHLEN, R. Thermodynamic properties of saturated and compressed liquid *n*-pentane 283

LANCASTER, N. M.; WORMALD, C. J. The excess molar enthalpies of $\{xH_2O+(1-x)C_2H_6\}$(g) and $\{xH_2O+(1-x)C_2H_4\}$(g) 295

Number 4, April 1985

MCGLASHAN, M. L. Phase equilibria in fluid mixtures. (The fifth Rossini Lecture of the Commission on Thermodynamics of IUPAC delivered on 14 August 1984 at its Conference on Chemical Thermodynamics held at McMaster University, Hamilton, Ontario, Canada) 301

ORTEGA, J.; PEÑA, J. A.; PAZ ANDRADE, M. I.; PINTOS, M.; ROMANI, L. Excess volumes of (benzene + each of several isomers of hexanol) at 298.15 K . 321

TAMURA, K.; MURAKAMI, S.; DOI, S. Speeds of sound, densities, and isentropic compressibilities of $\{xc\text{-}C_6H_{12} + (1-x)C_6H_5CH_3\}$, $\{xc\text{-}C_6H_{11}CH_3 + (1-x)C_6H_6\}$, and $\{xc\text{-}C_6H_{11}CH_3 + (1-x)C_6H_5CH_3\}$, from 293.15 to 303.15 K 325

CHRISTENSEN, J. J.; CORDRAY, D. R.; ZEBOLSKY, D. M.; IZATT, R. M. The excess enthalpies of (propane + dichlorodifluoromethane) from 353.15 to 398.15 K at 4.45 MPa 335

O'HARE, P. A. G. Thermochemistry of inorganic sulfur compounds. IV. Standard molar enthalpy of formation of sulfur hexafluoride . . . 349

LONGHI, P.; MUSSINI, T.; PENOTTI, F.; RONDININI, S. Thermodynamics of the cell $H_2|HCl|AgCl|Ag$ in (acetonitrile + water) solvents 355

MAJER, V.; SVOBODA, V.; LENCKA, M. Enthalpies of vaporization and cohesive energies of dimethylpyridines and trimethylpyridines 365

DIEZ, D.; RUIZ, B.; ROYO, F. M.; GUTIÉRREZ LOSA, C. Excess molar enthalpies at 298.15 K of (cyclohexane + a methylpyridine) and of (benzene + a methylpyridine) 371

CHHOR, K.; BOCQUET, J. F.; POMMIER, C. Low-temperature adiabatic calorimeter with an automatic data-acquisition system for the temperature range 10 to 310 K. The molar heat capacity of $NH_3(CH_2)_3NH_3MnCl_4$ 379

O'HARE, P. A. G.; JOHNSON, G. K. Rubidium chromate: standard molar enthalpies of solution and formation and the standard molar entropy at 298.15 K, enthalpy increments relative to 298.15 K, the high-temperature heat capacity, and temperatures, enthalpies, and entropies of transition and melting. Thermodynamic properties of Rb_2CrO_4 from 298.15 to 1500 K 391

Number 5, May 1985

BOERIO-GOATES, J.; LOPEZ DE LA FUENTE, F. L.; CHEDA, J. A. R.; WESTRUM, E. F., JR. Thermodynamics of thallium alkanoates. I. Heat capacity and thermodynamic functions of thallium(I) *n*-hexanoate 401

NGEYI, S. P.; LOPEZ DE LA FUENTE, F. L.; CHEDA, J. A. R.; FERNANDEZ-MARTIN, F.; WESTRUM, E. F., JR. Thermodynamics of thallium alkanoates. II. Heat capacity and thermodynamic functions of thallium(I) *n*-heptanoate 409

OTÍN, S.; MUÑOZ, J.; TOLEDO, M. J.; VELASCO, I.; GUTIÉRREZ LOSA, C. Excess molar enthalpies of (a halogenoaromatic compound + toluene) . . 419

HILPERT, K.; VISWANATHAN, R.; GINGERICH, K. A.; GERADS, H.; KOBERTZ, D. Vaporization of $FeI_2(s)$ and the thermochemistry of $FeI_2(g)$, $(FeI_2)_2(g)$, $(FeI_2)_3(g)$, and $FeI_3(g)$ 423

WORMALD, C. J.; COLLING, C. N. Excess enthalpies of (water + hydrogen)(g) up to 698.2 K and 11.13 MPa 437

NAVARRO, J. M.; BALOIRA, M.; BRAVO, R.; PAZ ANDRADE, M. I.; PINTOS, M. Excess enthalpies of (*n*-alkanol + ethyl ethanoate or ethyl propanoate) at 298.15 K 447

LETCHER, T. M.; HEYWARD, C.; PERKINS, D. M. Excess enthalpies and excess volumes of (cyclohexylbenzene + an *n*-alkane) at 298.15 K 453

LETCHER, T. M. The excess enthalpies and excess volumes of (cyclodecane + a bicyclic compound) at 298.15 K 459

CORDFUNKE, E. H. P.; OUWELTJES, W. Standard enthalpies of formation of uranium compounds. XII. Anhydrous phosphates 465

O'HARE, P. A. G.; BASILE, L.; APPELMAN, E. H. Thermochemistry of inorganic sulfur compounds. V. Peroxymonosulfate revisited: standard molar enthalpies of formation of $KHSO_5 \cdot H_2O$(cr), $KHSO_5$(cr), and HSO_5^-(aq) 473

ROBIE, R. A.; WIGGINS, L. B.; BARTON, P. B., JR.; HEMINGWAY, B. S. Low-temperature heat capacity and entropy of chalcopyrite ($CuFeS_2$): estimates of the standard molar enthalpy and Gibbs free energy of formation of chalcopyrite and bornite (Cu_5FeS_4) 481

MOLLERUP, J. Measurement of the volumetric properties of methane and ethene at 310 K at pressures to 70 MPa and of propene from 270 to 345 K at pressures to 3 MPa by the Burnett method 489

Number 6, June 1985

BENSON, G. C.; D'ARCY, P. J.; KUMARAN, M. K. Excess molar isobaric heat capacities of (2-methylpentan-1-ol + *n*-hexane) and (2-ethylbutan-1-ol + *n*-hexane) at 298.15 K 501

STEELE, W. V.; GAMMON, B. E.; SMITH, N. K.; CHICKOS, J. S.; GREENBERG, A.; LIEBMAN, J. F. The standard molar enthalpy of formation of 2,3-diphenylcycloprop-2-en-1-one 505

EWING, M. B.; JOHNSON, K. A.; MCGLASHAN, M. L. The liquid–liquid critical state of (cyclohexane + methanol). I. Excess molar enthalpies for $\{(1-x)c\text{-}C_6H_{12} + xCH_3OH\}$ within 1.3 K of the critical temperature . 513

HUTCHINGS, R. S.; VAN HOOK, W. A. Excess molar volumes and deviations from congruence of some binary solutions of an *n*-alkane in another *n*-alkane 523

HUTCHINGS, R. S.; VAN HOOK, W. A. Excess molar Gibbs free energies of pairs of *n*-alkanes, the two *n*-alkanes differing by one or two carbon atoms 531

SAITO, K.; ATAKE, T.; CHIHARA, H. Molar heat capacity and thermodynamic properties of *p*-quaterphenyl 539

EWING, M. B.; MCGLASHAN, M. L.; TRUSLER, J. P. M. The speed of sound in gases. I. A cylindrical interferometer and the speed of sound in argon and in krypton 549

GENS, R.; FUGER, J.; MORSS, L. R.; WILLIAMS, C. W. Thermodynamics of actinide perovskite-type oxides. III. Molar enthalpies of formation of Ba_2MAnO_6 (M = Mg, Ca, or Sr; An = U, Np, or Pu) and of M_3PuO_6 (M = Ba or Sr) 561

MANZUROLA, E.; APELBLAT, A. Apparent molar volumes of uranyl nitrate, uranyl chloride, and uranyl sulphate in water at 298.15 K 575

MANZUROLA, E.; APELBLAT, A. Apparent molar volumes of citric, tartaric, malic, succinic, maleic, and acetic acids in water at 298.15 K . . 579

RAGHAVAN, S.; IYENGAR, G. N. K.; ABRAHAM, K. P. Determination of the thermodynamic properties of $\{xMgO+(1-x)MnO\}$(s, sln) from a solid-electrolyte galvanic cell in the temperature range 1163 to 1318 K . . 585

O'HARE, P. A. G.; JOHNSON, G. K. Thermochemistry of inorganic sulfur compounds. VI. Standard molar enthalpy of formation at 298.15 K and the high-temperature thermodynamic properties of $NaLiSO_4$; implications of the results for $(0.5Na_2SO_4+0.5Li_2SO_4)$(l) . . . 593

Number 7, July 1985

IWASE, M.; SUGINO, S.; ICHISE, E.; WASEDA, Y. Application of Na^+-ion-conducting β-Al_2O_3 to thermodynamic studies of liquid sodium-based alloys: $\{xNa+(1-x)Cd\}$, $\{xNa+(1-x)In\}$, and $\{xNa+(1-x)Sn\}$. 601

O'HARE, P. A. G. Thermochemistry of uranium compounds. XV. Calorimetric measurements on UCl_4, UO_2Cl_2, and UO_2F_2, and the standard molar enthalpy of formation at 298.15 K of UCl_4 611

SCAIFE, W. G. S.; LYONS, C. G. R. Dielectric behaviour of freezing *n*-decane and *n*-pentadecane 623

AWWAD, A. M.; ALLOS, E. I.; KANBOUR, F. I. Thermodynamic properties of binary mixtures containing *N*-methylpyrrolidone. I. Excess molar volumes of (an *n*-alkanol + *N*-methylpyrrolidone) 635

PIMERZIN, A. A.; NESTEROVA, T. N.; ROZHNOV, A. M. Equilibria of isomeric transformations and relations between thermodynamic properties of secondary alkylbenzenes 641

NESTEROVA, T. N.; ROZHNOV, A. M.; MALOVA, T. N.; KOVZEL, E. N. Molar enthalpies of formation of isopropylchlorobenzenes derived from equilibrium measurements 649

HALL, P. G.; ARMITAGE, D. A.; LINFORD, R. G. Heat capacity of silver tungstate between 80 and 500 K 657

BOERIO-GOATES, J.; GOATES, S. R.; OTT, J. B.; GOATES, J. R. Enthalpies of formation of molecular addition compounds in tetrachloromethane + *p*-xylene, + toluene, and + benzene from (solid + liquid) phase equilibria 665

BRUNNER, E. Fluid mixtures at high pressures. I. Phase separation and critical phenomena of 10 binary mixtures of (a gas + methanol) . . . 671

ARMITAGE, D. A.; DUFFIELD, A.; HILL, R. J.; SWINTON, F. L. Excess volumes of (carbon disulphide + *n*-hexane) 681

BOUWSTRA, J. A. LEEUW, V. V. DE; MILTENBURG, J. C. VAN. Properties of mixed-crystalline organic material prepared by zone levelling. IV. Melting properties and excess enthalpies of (*trans*-azobenzene + *trans*-stilbene) . 685

NOTES

LOZANO, J. M.; OTIN, S.; GUTIÉRREZ LOSA, C. Excess enthalpies for (1-chlorohexane + a diketone) 697

KUMARAN, M. K.; BENSON, G. C. Excess volumes of (decan-1-ol + 2,2,3-trimethylbutane) 699

LINFORD, R. G.; THORNHILL, D. G. T. Solubilities of a gas in a crown ether 701

Number 8, August 1985

SIPOWSKA, J. T.; WIECZOREK, S. A. Excess molar volumes of (cyclohexanol + *n*-heptane) at 303.15 and 308.15 K 703

LETCHER, T. M.; HEYWARD, C. The excess enthalpies and excess volumes of (cyclohexylbenzene + a cycloalkane) at 298.15 K 707

PÉREZ, P.; ROYO, F.; GRACIA, M.; GUTIÉRREZ LOSA, C. H_m^E and V_m^E of some (1-chlorobutane + alkanol or cyclohexanol) mixtures 711

BATTLER, J. R.; ROWLEY, R. L. Excess enthalpies between 293 and 323 K for constituent binaries of ternary mixtures exhibiting partial miscibility . 719

MORON, M. C.; PÉREZ, P.; GRACIA, M.; GUTIÉRREZ LOSA, C. Excess molar enthalpies of mixtures containing 1,2-dibromoethane 733

GREENBERG, J. H.; GUSKOV, V. N.; LAZAREV, V. B. Vapour-pressure investigation of thermodynamics of non-stoichiometric crystals. Sublimation of β-Zn_3As_2 739

NAGATA, I.; TAMURA, K. Excess molar enthalpies of (propan-1-ol + ethyl ethanoate + cyclohexane) and (propan-2-ol + ethyl ethanoate + cyclohexane) at 298.15 K 747

JOHNSON, G. K.; MURRAY, W. T.; VAN DEVENTER, E. H.; FLOTOW, H. E. The thermodynamic properties of zirconium ditelluride $ZrTe_2$ to 1500 K . 751

FUJIHARA, I.; OGAWA, H.; MURAKAMI, S. Excess volumes of *cis*-decalin + benzene, + toluene, + isooctane, and + *n*-heptane at 298.15 K . . 761

JACKOWSKI, A. W. Liquid-vapour equilibria in $\{xCH_2Cl_2 + (1-x)C_4H_9CH : CH_2\}$ at 298.17 K 765

APELBLAT, A. Enthalpy of solution of lithium sulfate and lithium sulfate monohydrate in water at 298.15 K 769

HALSTEAD, K. E.; SEDDON, D. J.; STAVELEY, L. A. K.; WEIR, R. D. The heat capacity of manganese and zinc hexafluorosilicates at low temperatures . 775

WALTERS, S. T.; STAVELEY, L. A. K. The heat capacity of potassium hexafluoroantimonate(V) from 21 to 284 K 781

CHRISTENSEN, J. J.; ZEBOLSKY, D. M.; IZATT, R. M. The excess enthalpies of (carbon dioxide + pyridine) at 470.15 and 573.15 K from 7.50 to 12.50 MPa 785

CROUCH-BAKER, S.; DICKENS, P. G.; KAY, S. A. Standard molar enthalpies of formation of $WO_3 \cdot 2H_2O$ and $WO_3 \cdot H_2O$ by solution calorimetry . 797

Number 9, September 1985

GLASER, M.; PETERS, C. J.; KOOI, H. J. VAN DER; LICHTENTHALER, R. N. Phase equilibria of (methane + *n*-hexadecane) and (p, V_m, T) of *n*-hexadecane . 803

RODRIGUEZ-NUÑEZ, E.; PAZ-ANDRADE, M. I.; BRAVO, R.; ORTEGA, J. Thermodynamic properties of (a xylene + an *n*-alkanol). III. Excess molar volumes at 298.15 K for 1-hexanol, 1-heptanol, and 1-octanol . . . 817

SANAHUJA, A.; CESARI, E. Enthalpy of solution of aqueous sodium chloride at 303.15, 308.15, and 313.15 K 823

KALINOWSKA, B.; WÓYCICKI, W. Heat capacities and excess heat capacities of (an alcohol + an unsaturated hydrocarbon). I. (Propan-1-ol + *n*-hex-1-ene) 829

MARSH, K. N.; ORGAN, P. P. Excess molar enthalpies and excess molar volumes for three- and four-component *n*-alkane mixtures simulating (*n*-hexane + *n*-hexadecane) 835

ROYO, F. M.; GUTIÉRREZ LOSA, C. Vapour pressures of (*n*-hexane + an α,ω-dichloroalkane) 843

NIKOLIĆ, A. D.; PERIŠIĆ-JANJIĆ, N. U.; PETROVIĆ, S. D.; KOBILAROV, N. L. A thermodynamic study of *N*-*n*-butylbenzamide solutions using gas-liquid chromatography 849

ROBINSON, D. S.; KIDNAY, A. J.; YESAVAGE, V. F. The enthalpy of *trans*-decalin between 360 and 700 K and at pressures to 10.3 MPa . . . 855

JACKOWSKI, A. W. Vapour pressure of (*n*-hexane + *n*-butyl octadecanoate). I. Isotropic liquid-vapour and smectic-isotropic liquid-vapour equilibria . 865

BRUNNER, E. Fluid mixtures at high pressures. II. Phase separation and critical phenomena of (ethane + an *n*-alkanol) and of (ethene + methanol) and (propane + methanol) 871

MACHÁT, V.; BOUBLÍK, T. Vapour-liquid equilibrium and excess molar volume of (*trans*-1,2-dichloroethene + cyclopentane) at 293.15 K . . . 887

SINKE, G. C.; MOSSNER, E. H.; CURNUTT, J. L. Enthalpies of solution and solubilities of calcium chloride and its lower hydrates 893

NOTE

UEDAIRA, HATSUHO; UEDAIRA, HISASHI. The relation between partial molar heat capacities and the number of equatorial hydroxyl groups of saccharides in aqueous solution 901

Number 10, October 1985

WORMALD, C. J.; LANCASTER, N. M. Excess molar enthalpies for (water + benzene)(g) and (water + cyclohexane)(g) 903

TOUHARA, H.; NAKANISHI, K. Excess molar enthalpies of methanol + pyridine, + methylpyridine, and + 2,6-dimethylpyridine . . . 909

ROGALSKI, M.; BYLICKI, A. Vapour pressures and excess Gibbs energies of {pyridine + (methyl- or ethyl- or *n*-propyl- or *n*-butyl-benzene)} at $T = 373.15$ K 915

ROGALSKI, M. Vapour pressures and excess Gibbs energies of {(3-methylpyridine or 3-ethylpyridine) + (methyl- or ethyl- or propyl- or butylbenzene)} at $T = 373.15$ K 921

DOWNIE, D. B.; MARTIN, J. F. Heat capacity of gamma-phase alloys of (silver + zinc) and (copper + zinc) 927

OTIN, S.; TOLEDO, M. J.; MUÑOZ, J.; VELASCO, I.; GUTIÉRREZ LOSA, C. Excess molar enthalpies of (a halogenobenzene + benzene or + toluene) and of some (a dihalogenobenzene + benzene) 935

ZAWISZA, A. High-pressure liquid-vapour equilibria, critical state, and $p(V_m, T, x)$ to 448.15 K and 4.053 MPa for $\{xC_6H_{14} + (1-x)CH_3OH\}$. 941

FIGUIÈRE, P.; SZWARC, H.; OGUNI, M.; SUGA, H. Calorimetric study of thiophene from 13 to 300 K. Emergence of two glassy crystalline states . 949

ARNTZ, H.; GOTTLIEB, K. High-pressure heat-flow calorimeter. Determination of the enthalpy of reaction for the synthesis of methyl *t*-butyl ether from methanol and 2-methylpropene 967

HAMAM, S. E. M.; BENSON, G. C.; KUMARAN, M. K. Excess enthalpies of (water + diethylene glycol) and (water + triethylene glycol) . . . 973

KUMAGAI, A.; TAKAHASHI, S. (Pressure, volume, temperature) behaviour of liquid 1,1-dichloroethane and 1,2-dichloroethane 977

GARCIA-SANCHEZ, F.; TREJO, A. Vapour pressure and critical constants of 1,2-dichloroethane 981

BRUNNER, E. Fluid mixtures at high pressures. III. Isothermal phase equilibria of (ethene + methanol) at 284.15 and 298.15 K 985

MARSH, K. N.; MÅNSSON, M. Standard molar enthalpies of formation of triethoxymethane and tetraethoxymethane by rotating bomb calorimetry 995

Number 11, November 1985

TOPOR, L.; KLEPPA, O. J. Enthalpies of formation of first-row transition-metal diborides by a new calorimetric method 1003

NAGATA, I. Isothermal (vapour + liquid) equilibria of propan-1-ol + acetonitrile and + trichloromethane and of (propan-1-ol + acetonitrile + trichloromethane) 1017

ABELLO, L.; CHHOR, K.; POMMIER, C. Thermodynamic studies on the successive phase transitions in $LiKSO_4$ crystals at low temperatures . 1023

CORDFUNKE, E. H. P.; MUIS, R. P.; WIJBENGA, G.; BURRIEL, R.; TO, M. (W. K.); ZAINEL, H.; WESTRUM, E. F., JR. Thermodynamics of uranium intermetallic compounds. I. Heat capacities of URu_3 and URh_3 from 5 to 850 K 1035

ROGALSKI, M. Vapour pressures and excess Gibbs energies of (2-methylpyridine + methylbenzene), (2,4-dimethylpyridine + 1,2-dimethylbenzene), and (2,6-dimethylpyridine + 1,2- or 1,3- or 1,4-dimethylbenzene or ethylbenzene) at $T = 373.15$ K 1045

ROGALSKI, M. Vapour pressures and excess Gibbs energies of (2,4,6-trimethylpyridine + 1,2,4- or 1,3,5-trimethylbenzene or *n*-propylbenzene or isopropylbenzene or *n*-butylbenzene or isobutylbenzene) at $T = 373.15$ K 1051

TAKAGI, T.; TERANISHI, H. Ultrasonic speeds and thermodynamics for (toluene + *o*-xylene) and (toluene + aniline) under high pressures 1057

SANAHUJA, A. Enthalpy of solution of KCl in water at 308.15 and 313.15 K 1063

PEREZ, P.; GRACIA, M.; GUTIÉRREZ LOSA, C. H_m^E and V_m^E for (2,2-dimethylbutane or 2-chloro-2-methylpropane + a monoalcohol) . . . 1067

AWWAD, A. M.; AL-MADAFI, S. F.; JBARA, K. A.; ALCHY, W. T. Excess molar volumes of (*N*-formylmorpholine + a glycol) at 298.15, 308.15, and 318.15 K 1073

CORDFUNKE, E. H. P.; CLUISTRA, R.; MILTENBURG, J. C. VAN. The thermodynamic properties of six compounds in (tellurium + oxygen + hydrogen) from 10 to 1000 K 1079

COLOMINA, M.; JIMÉNEZ, P.; ROUX, M. V.; TURRIÓN, C. Thermochemical properties of benzoic acid derivatives. XII. Vapour pressures and enthalpies of sublimation and formation of five dimethoxybenzoic acids 1091

LUO, C.-W.; WEBER, J. H.; TAO, L. C. Vapor–liquid equilibria of (methanol + methylcyclopentane) 1097

Number 12, December 1985

NAGATA, I.; NAKAMURA, S. Liquid–liquid equilibria of (cyclohexane + acetonitrile + methylcyclohexane + toluene) and of {(acetonitrile + methylcyclohexane) + benzene or +toluene or +cyclohexane or +chlorobenzene} at 298.15 K 1103

SHINSAKA, K.; GEE, N.; FREEMAN, G. R. Densities against temperature of 17 organic liquids and of solid 2,2-dimethylpropane 1111

AZNAR, E.; RUIZ, B.; GUTIÉRREZ LOSA, C. Excess molar enthalpies at 298.15 K of (cyclohexane + a methylquinoline) and of (benzene + a methylquinoline) 1121

ORTEGA, J.; MATOS, J. S.; PAZ ANDRADE, M. I.; JIMENEZ, E. Excess molar volumes of (ethyl formate or ethyl acetate + an isomer of hexanol) at 298.15 K 1127

COLINET, C.; PASTUREL, A.; BUSCHOW, K. H. J. Molar enthalpies of formation of $LnAl_2$ compounds 1133

KOHLEN, R.; KRATZKE, H.; MÜLLER, S. Thermodynamic properties of saturated and compressed liquid difluorochloromethane . . . 1141

LAINEZ, A.; WILHELM, E.; ROUX-DESGRANGES, G.; GROLIER, J.-P. E. Excess molar quantities of (a halogenated *n*-alkane + an *n*-alkane). A comparative study of mixtures containing either 1-chlorobutane or 1,4-dichlorobutane. 1153

VEGA, C. A.; MUÑIZ, M. DE LOS A. Standard potential of the (Ag + AgCl) electrode in (acetonitrile + water) 1163

WORMALD, C. J.; YERLETT, T. K. A new enthalpy-increment calorimeter. Enthalpy increments for *n*-hexane 1171

GIRARD, C.; BARET, R.; MIANE, J. M.; BROS, J. P. Excess molar enthalpies of $\{(1-x_2-x_3)Al + x_2Bi + x_3Ga\}(l)$ alloys 1187

NOTE

ORTEGA, J.; PAZ ANDRADE, M. I.; BRAVO, R. Excess molar volumes at 298.15 K of (benzene + 2-methylpentan-3-ol) and of (benzene + 2-ethylbutan-1-ol) 1199

INDEX TO VOLUME 17 1201